

## **UNIT-2: Mass Spectrometry**

Basic principles, Instrumentation - Mass spectrometer, electron Ionization (Electron Impact ionization, EI), Molecular ions, metastable ions, Isotope abundance. Basic fragmentation types. Fragmentation patterns in Toluene, 2-Butanol, Butanaldehyde, Propionic acid.

### **Essay Question 1:**

**Discuss the basic principles of Mass Spectrometry. Explain in detail the instrumentation involved in mass spectrometry with a focus on Electron Ionization (EI).**

#### **Mass Spectrometry Principles:**

*Mass spectrometry (MS) is an analytical technique used to identify chemical substances based on the mass-to-charge ratio ( $m/z$ ) of their ions. The basic principles of mass spectrometry include:*

- 1. Ionization: The sample is ionized to convert neutral molecules into charged ions.*
- 2. Mass Analysis: The ions are separated based on their mass-to-charge ( $m/z$ ) ratio by passing through an electric or magnetic field.*
- 3. Detection: The separated ions are detected, and a mass spectrum is generated, showing the abundance of ions at each  $m/z$  value.*

*The mass spectrum provides information about the molecular weight and structure of the compound being analyzed.*

#### **Instrumentation of Mass Spectrometry:**

*A typical mass spectrometer consists of the following components:*

- 1. Sample Inlet: The sample is introduced into the ion source.*
- 2. Ion Source: This is where ionization occurs. In Electron Ionization (EI), the sample molecules are bombarded with high-energy electrons (typically 70 eV), causing them to lose an electron and form a positive ion, often referred to as the molecular ion ( $M^+$ ).*
- 3. Mass Analyzer: The ions are accelerated into a mass analyzer where they are separated based on their  $m/z$  ratio. Common types of mass analyzers include quadrupoles, time-of-flight (TOF), and magnetic sectors.*


4. *Detector: The ions are detected as they reach the detector, where they generate an electrical signal proportional to their abundance.*
5. *Data System: The output from the detector is processed to generate the mass spectrum, a plot of ion abundance versus  $m/z$  ratio.*

#### *Electron Ionization (EI):*


*In EI, high-energy electrons collide with neutral molecules, causing ionization by removing an electron from the molecule. The molecular ion ( $M^+$ ) formed is usually unstable and fragments into smaller ions. The fragmentation pattern provides valuable structural information about the compound.*

- *Advantages of EI: Provides reproducible spectra and extensive fragmentation patterns, which help in structure elucidation.*
- *Limitations: It can result in excessive fragmentation, making it difficult to detect the molecular ion for certain compounds.*

## Mass spectrometry



# MASS SPECTROMETRY



## Essay Question 2:

Explain in detail the fragmentation patterns observed in 1. Toluene, 2. Butanol-2, 3. Buteraldehyde and 4. Propanoic Acid in mass spectrometry.

### 1. Toluene ( $C_6H_5CH_3$ ):

- **Molecular ion ( $M^+$ ):** The molecular ion for toluene appears at  $m/z = 92$ .
- **Fragmentation:** The major fragmentation involves the loss of a methyl group ( $CH_3$ , 15 amu), resulting in a benzyl ion ( $C_6H_5^+$ ) at  $m/z = 77$ . Other common fragments include tropylium ion ( $C_7H_7^+$ ) at  $m/z = 91$ , which forms via rearrangement.

### 2. 2-Butanol ( $C_4H_9OH$ ):

- **Molecular ion ( $M^+$ ):** The molecular ion for 2-butanol appears at  $m/z = 74$ .
- **Fragmentation:** The common fragmentation involves the loss of a water molecule ( $H_2O$ , 18 amu), resulting in a fragment at  $m/z = 56$ . Other fragmentation patterns include cleavage between the carbon-carbon bonds, yielding ions like  $C_2H_5^+$  at  $m/z = 29$  and  $CH_3^+$  at  $m/z = 15$ .

### 3. Buteraldehyde ( $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-CHO}$ ):

Molecular formula:  $\text{C}_4\text{H}_8\text{O}$

Molecular weight: 72 amu

Structure:  $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-CHO}$

#### Key Fragmentation Patterns:

Peak (m/z)	Fragment	Explanation
72 ( $\text{M}^+$ )	Molecular ion	Weak due to instability of aldehyde cation
57	$[\text{M} - \text{CH}_3]^+$	Loss of a methyl group (-15), forms $\text{CH}_3\text{-CH}_2\text{-CH=O}^+$ (stable acylium ion)
44	$[\text{CH}_3\text{-CH}_2\text{-CO}]^+$ or $\text{CH}_2\text{CHO}^+$	Cleavage between $\alpha$ and $\beta$ carbon ( $\alpha$ -cleavage) to give resonance-stabilized ion
43	$[\text{CH}_3\text{CO}]^+$ or $\text{CH}_3\text{CH=O}^+$	Acylium ion, very stable; commonly observed in aldehydes
29	$[\text{CH}_3\text{CH}_2]^+$	Ethyl carbocation, from deeper fragmentation

#### Mechanism Highlights:

- **$\alpha$ -Cleavage** (common in carbonyl compounds): Breaks the bond adjacent to the carbonyl group to generate resonance-stabilized cations like  $\text{CH}_3\text{CH=O}^+$ .
- **McLafferty Rearrangement** is not prominent in straight-chain aliphatic aldehydes like butanal due to lack of  $\gamma$ -hydrogens required for the rearrangement.

### 4. Propanoic Acid

#### Short Answer Question 1:

*What is the nitrogen rule in mass spectrometry, and how is it useful in determining molecular structures?*

*The nitrogen rule in mass spectrometry states that organic compounds containing an odd number of nitrogen atoms will have an odd molecular ion mass ( $\text{M}^+$ ), while compounds with an even number of nitrogen atoms or no nitrogen will have an even molecular ion mass.*

### *Utility in Structural Determination:*

- *This rule helps in quickly determining the presence of nitrogen in a compound.*
- *If the molecular ion has an odd  $m/z$  value, it suggests that the molecule contains an odd number of nitrogen atoms.*
- *This is particularly useful when analyzing complex molecular structures to predict elemental composition.*

### *Short Answer Question 2:*

*Describe the McLafferty rearrangement and its significance in mass spectrometry.*

*The McLafferty rearrangement is a fragmentation pattern observed in mass spectrometry involving a six-membered ring transition state. It typically occurs in compounds with a carbonyl group (C=O) and a gamma-hydrogen (a hydrogen atom located three carbon atoms away from the carbonyl carbon).*

### *Mechanism:*

- *In this rearrangement, a hydrogen atom from the gamma position is transferred to the carbonyl oxygen, resulting in the formation of a double bond between the alpha and beta carbon atoms and cleavage between the alpha and beta positions.*
- *This generates a neutral molecule and a radical cation fragment.*

### *Significance:*

- *The McLafferty rearrangement provides specific structural information, especially in carbonyl-containing compounds, as it indicates the presence of a gamma-hydrogen.*

### *Short Answer Question 3:*

*Define isotopic abundance and explain its significance in mass spectrometry analysis.*

*Isotopic abundance refers to the relative percentage of different isotopes of an element found in nature. In mass spectrometry, isotopes produce peaks at different  $m/z$  values corresponding to their masses.*

### *Significance:*

- *Isotopic abundance is important for identifying elements with naturally occurring isotopes (e.g., chlorine, bromine). The presence of isotopic peaks (e.g.,  $M^+$  and  $M^+2$ ) in the*

*mass spectrum helps in distinguishing between different elements and confirming molecular composition.*

- *For example, chlorine has two common isotopes,  $^{35}\text{Cl}$  and  $^{37}\text{Cl}$ , which result in a characteristic M and M+2 peak ratio of 3:1, aiding in compound identification.*

*Short Answer Question 4:*

*What are metastable ions? Describe their formation and characteristics.*

*Metastable ions are ions that have insufficient internal energy to fragment immediately after formation, but they undergo fragmentation as they pass through the mass spectrometer.*

*Formation:*

- *These ions form during the transition between the ion source and the detector in a mass spectrometer.*
- *Metastable ions fragment in regions of the spectrometer where no electric or magnetic fields are present (often in the drift region).*

*Characteristics:*

- *They produce broad, diffuse peaks in the mass spectrum called metastable peaks.*
- *These peaks appear at non-integer  $m/z$  values and provide information about the fragmentation pathways of the parent ion, helping in structural elucidation.*